

11.1.2 *d*-Separation without Tears (Chapter 1, pp. 16–18)

At the request of many who have had difficulties switching from algebraic to graphical thinking, I am including a gentle introduction to *d*-separation, supplementing the formal definition given in Chapter 1, pp. 16–18.

Introduction

d-separation is a criterion for deciding, from a given causal graph, whether a set X of variables is independent of another set Y , given a third set Z . The idea is to associate “dependence” with “connectedness” (i.e., the existence of a connecting path) and “independence” with “unconnectedness” or “separation.” The only twist on this simple idea is to define what we mean by “connecting path,” given that we are dealing with a system of directed arrows in which some vertices (those residing in Z) correspond to measured variables, whose values are known precisely. To account for the orientations of the arrows we use the terms “*d*-separated” and “*d*-connected” (*d* connotes “directional”). We start by considering separation between two singleton variables, x and y ; the extension to sets of variables is straightforward (i.e., two sets are separated if and only if each element in one set is separated from every element in the other).

Unconditional Separation

Rule 1: x and y are *d*-connected if there is an unblocked path between them.

By a “path” we mean any consecutive sequence of edges, disregarding their directionalities. By “unblocked path” we mean a path that can be traced without traversing a pair of arrows that collide “head-to-head.” In other words, arrows that meet head-to-head do not constitute a connection for the purpose of passing information; such a meeting will be called a “collider.”

Example 11.1.1 The graph in Figure 11.1 contains one collider, at t . The path $x - r - s - t$ is unblocked, hence x and t are *d*-connected. So also is the path $t - u - v - y$, hence t and y are *d*-connected, as well as the pairs u and y , t and v , t and u , x and s , etc. However, x and y are not *d*-connected; there is no way of tracing a path from x to y without traversing the collider at t . Therefore, we conclude that x and y are *d*-separated, as well as x and v , s and u , r and u , etc. (In linear models, the ramification is that the covariance terms corresponding to these pairs of variables will be zero, for every choice of model parameters.)




Figure 11.2 The set $Z = \{r, v\}$ d -separates x from t and t from y .




Figure 11.3 s and y are d -connected given p , a descendant of the collider t .

Blocking by Conditioning

Motivation: When we measure a set Z of variables, and take their values as given, the conditional distribution of the remaining variables changes character; some dependent variables become independent, and some independent variables become dependent. To represent this dynamic in the graph, we need the notion of “conditional d -connectedness” or, more concretely, “ d -connectedness, conditioned on a set Z of measurements.”

Rule 2: x and y are d -connected, conditioned on a set Z of nodes, if there is a collider-free path between x and y that traverses no member of Z . If no such path exists, we say that x and y are d -separated by Z . We also say then that every path between x and y is “blocked” by Z .

Example 11.1.2 Let Z be the set $\{r, v\}$ (marked by circles in Figure 11.2). Rule 2 tells us that x and y are d -separated by Z , and so also are x and s , u and y , s and u , etc. The path $x - r - s$ is blocked by Z , and so also are the paths $u - v - y$ and $s - t - u$. The only pairs of unmeasured nodes that remain d -connected in this example, conditioned on Z , are s and t and u and t . Note that, although t is not in Z , the path $s - t - u$ is nevertheless blocked by Z , since t is a collider, and is blocked by Rule 1.

Conditioning on Colliders

Motivation: When we measure a common effect of two independent causes, the causes become dependent, because finding the truth of one makes the other less likely (or “explained away,” p. 17), and refuting one implies the truth of the other. This phenomenon (known as Berkson paradox, or “explaining away”) requires a slightly special treatment when we condition on colliders (representing common effects) or on any descendant of a collider (representing evidence for a common effect).

Rule 3: If a collider is a member of the conditioning set Z , or has a descendant in Z , then it no longer blocks any path that traces this collider.

Example 11.1.3 Let Z be the set $\{r, p\}$ (again, marked with circles in Figure 11.3). Rule 3 tells us that s and y are d -connected by Z , because the collider at t has a descendant (p) in Z , which unblocks the path $s - t - u - v - y$. However, x and u are still d -separated by Z , because although the linkage at t is unblocked, the one at r is blocked by Rule 2 (since r is in Z).

This completes the definition of d -separation, and readers are invited to try it on some more intricate graphs, such as those shown in Chapter 1, Figure 1.3.

Typical application: Consider Example 11.1.3. Suppose we form the regression of y on p , r , and x ,

$$y = c_1p + c_2r + c_3x + \epsilon,$$

and wish to predict which coefficient in this regression is zero. From the discussion above we can conclude immediately that c_3 is zero, because y and x are d -separated given p and r , hence y is independent of x given p and r , or, x cannot offer any information about y once we know p and r . (Formally, the partial correlation between y and x , conditioned on p and r , must vanish.) c_1 and c_2 , on the other hand, will in general not be zero, as can be seen from the graph: $Z = \{r, x\}$ does not d -separate y from p , and $Z = \{p, x\}$ does not d -separate y from r .

Remark on correlated errors: Correlated exogenous variables (or error terms) need no special treatment. These are represented by bi-directed arcs (double-arrowed), and their arrowheads are treated as any other arrowheads for the purpose of path tracing. For example, if we add to the graph in Figure 11.3 a bi-directed arc between x and t , then y and x will no longer be d -separated (by $Z = \{r, p\}$), because the path $x - t - u - v - y$ is d -connected – the collider at t is unblocked by virtue of having a descendant, p , in Z .